

United States Department of Agriculture Natural Resources Conservation Service

Ecological Site Description

Site Type: Rangeland

Site Name: Saline Subirrigated (SS), 15-19" P.Z., Foothills and Mountains West

Site ID: R043BY242WY

Major Land Resource Area: 43B-Central Rocky Mountains

Physiographic Features

This site occurs on nearly level land along perennial or intermittent streams, and near seeps, sloughs, or springs. It is also found on broad, low lake terraces, lake plains, on alluvial bottoms, and poorly-drained bottom lands adjacent to stream channels. These areas receive additional run-in water from higher sites and from a fluctuating water table, well within the root zone.

Landform: alluvial fans, drainage ways & stream terraces

Aspect: N/A

	<u>Minimum</u>	<u>Maximum</u>	
Elevation (feet):	5600	8300	
Slope (percent):	0	10	(mostly <3%)
Water Table Depth (inches):	0	306	
Flooding:			
Frequency:	occasional	frequent	
Duration:	very brief	brief	
Ponding:			
Depth (inches):	0	0	
Frequency:	none	none	
Duration:	none	none	
Runoff Class:	negligible	high	

Climatic Features

Annual precipitation ranges from 15-19 inches per year. Wide fluctuations may occur in yearly precipitation and result in more dry years than those with more than normal precipitation. Temperatures show a wide range between summer and winter and between daily maximums and minimums. This is predominantly due to the high elevation and dry air, which permits rapid incoming and outgoing radiation. Cold air outbreaks in winter move rapidly from northwest to southeast and account for extreme minimum temperatures. Extreme storms may occur during the winter, but most severely affect ranch operations during late winter and spring.

Prevailing winds are from the southwest, and strong winds are less frequent than over other areas of Wyoming. Occasional storms, however, can bring brief periods of high winds with gusts exceeding 50 mph.

Growth of native cool season plants begins about May 15 and continues to about August 15.

The following information is from the "Jackson" climate station:

	<u>Minimum</u>	<u>Maximum</u>	<u>5 yrs. out of 10 between</u>
Frost-free period (days):	12	60	July 9 – August 12
Freeze-free period (days):	42	100	June 20 – August 26

Annual Precipitation (inches): <11.98 >19.69 (2 years in 10)

Mean annual precipitation: 17.00 inches

Mean annual air temperature: 38.9°F (23.3°F Avg. Min. to 54.5°F Avg. Max.)

For detailed information visit the Natural Resources Conservation Service National Water and Climate Center at <http://www.wcc.nrcs.usda.gov/cgibin/state.pl?state=wy> website. Other climate stations representative of this precipitation zone include "Afton" in Lincoln County; and "Darwin Ranch" in Teton County.

Influencing Water Features

<u>Wetland Description:</u>	<u>System</u>	<u>Subsystem</u>	<u>Class</u>	<u>Sub-class</u>
None	None	None	None	None

Stream Type: C (Rosgen)

Representative Soil Features

Soils of this site are mostly deep, somewhat poorly drained and affected by wetness and salt. Textures are range from loamy fine sand to clay, with loam to clay being most common. Salt crusts are commonly found on mounds during dry periods. The water table fluctuates during the growing season and is generally between 20 to 40 inches most years. The soil and soil water have enough salinity to restrict plant establishment and growth. Mottling or gleying may occur within 20 to 40 inches of the surface.

Major Soil Series correlated to this site include: Bear Lake Series

Parent Material Kind: alluvium

Parent Material Origin: mixed

Surface Texture: loam, silt loam, silty clay loam

Surface Texture Modifier: none

Subsurface Texture Group: loam, sandy loam, clay loam, loamy sand

Surface Fragments ≤ 3" (% Cover): 0-10

Surface Fragments > 3" (%Cover): 0

Subsurface Fragments ≤ 3" (% Volume): 0-30

Subsurface Fragments > 3" (% Volume): 0-10

	<u>Minimum</u>	<u>Maximum</u>
Drainage Class:	somewhat poorly	moderately well
Permeability Class:	moderately slow	moderate
Depth (inches):	20	>60
Electrical Conductivity (mmhos/cm) ≤20":	8	>16
Sodium Absorption Ratio ≤20":	0	10
Soil Reaction (1:1 Water) ≤20":	7.4	9.0
Soil Reaction (0.1M CaCl2) ≤20":	NA	NA
Available Water Capacity (inches) ≤30":	3	6
Calcium Carbonate Equivalent (percent) ≤20":	10	30


Plant Communities

Ecological Dynamics of the Site:

As this site deteriorates from improper grazing management, species such as inland saltgrass, alkali bluegrass, and alkali muhly increase. Grasses such as alkali sacaton, basin wildrye, and Nuttall's alkaligrass will decrease in frequency and production.

The Historic Climax Plant Community (description follows the plant community diagram) has been determined by study of rangeland relic areas, or areas protected from excessive disturbance. Trends in plant communities going from heavily grazed areas to lightly grazed areas, seasonal use pastures, and historical accounts have also been used.

The following is a State and Transition Model Diagram that illustrates the common plant communities (states) that can occur on the site and the transitions between these communities. The ecological processes will be discussed in more detail in the plant community narratives following the diagram.



BMA – Brush Management (all methods)
 BMC – Brush Management (chemical)
 BMF – Brush Management (fire)
 BMM – Brush Management (mechanical)
 CSP – Chemical Seedbed Preparation
 CSLG – Continuous Season-long Grazing
 DR – Drainage
 CSG – Continuous Spring Grazing
 HB – Heavy Browse
 HCSLG – Heavy Continuous Season-long Grazing
 HI – Heavy Inundation
 LPG – Long-term Prescribed Grazing
 MT – Mechanical Treatment (chiseling, ripping, pitting)

NF – No Fire
 NS – Natural Succession
 NWC – Noxious Weed Control
 NWI – Noxious Weed Invasion
 NU – Nonuse
 P&C – Plow & Crop (including hay)
 PG – Prescribed Grazing
 RPT – Re-plant Trees
 RS – Re-seed
 SGD – Severe Ground Disturbance
 SHC – Severe Hoof Compaction
 WD – Wildlife Damage (Beaver)
 WF – Wildfire

Plant Community Composition and Group Annual Production
Reference Plant Community (HCPC)

COMMON NAME/GROUP NAME	SCIENTIFIC NAME	SYMBOL	Annual Production (Normal Year)		
			Total: 3800		
			Group	lbs./acre	% Comp.
GRASSES AND GRASS-LIKES					
GRASSES/GRASSLIKES					
alkali sacaton	Sporobolus airoides	SPAI	1	1520 - 1900	40 - 50
basin wildrye	Leymus cinereus	LECI4	2	190 - 380	5 - 10
Nuttall's alkaligrass	Puccinellia nuttalliana	PUNU2	3	190 - 380	5 - 10
Alkali bluegrass	Poa juncifolia (syn. to P. secunda)	POJU (POSE)	4	190 - 380	5 - 10
inland saltgrass	Distichlis spicata	DISP	5	38 - 380	1 - 10
MISC. GRASSES/GRASSLIKES			6	190 - 570	5 - 15
alkali muhly	Muhlenbergia asperifolia	MUAS	6	0 - 190	0 - 5
Bearded wheatgrass	Elymus scribneri	ELSC4	6	0 - 190	0 - 5
bentgrass	Agrostis spp.	AGROS2	6	0 - 190	0 - 5
inland sedge	Carex interior	CAIN11	6	0 - 190	0 - 5
Little barley	Hordeum pusillum	HOPU	6	0 - 190	0 - 5
mat muhly	Muhlenbergia richardsonis	MURI	6	0 - 190	0 - 5
Nebraska sedge	Carex nebrascensis	CANE2	6	0 - 190	0 - 5
tufted hairgrass	Deschampsia caespitosa	DECA18	6	0 - 190	0 - 5
rhizomatous wheatgrass	Pascopyrum smithii	PASM	6	0 - 190	0 - 5
Rush	Juncus spp.	JUNCU	6	0 - 190	0 - 5
Sedge	Carex spp.	CAREX	6	0 - 190	0 - 5
Shortawn foxtail	Alopecurus aequalis	ALAE	6	0 - 190	0 - 5
other perennial grasses (native)		2GP	6	0 - 190	0 - 5
FORBS			7	38 - 380	1 - 10
arrowgrass	Triglochin spp.	TRIGL	7	0 - 190	0 - 5
Cinquefoil (herbaceous)	Potentilla spp.	POTEN	7	0 - 190	0 - 5
Elk thistle	Cirsium foliosum	CIFO	7	0 - 190	0 - 5
Gentian	Gentiana spp.	GENTI	7	0 - 190	0 - 5
milkvetch	Astragalus spp.	ASTRA	7	0 - 190	0 - 5
Mountain dandelion	Agoseris spp.	AGOSE	7	0 - 190	0 - 5
Owl's-clover	Orthocarpus spp.	ORTHO	7	0 - 190	0 - 5
other perennial forbs (native)		2FP	7	0 - 190	0 - 5
TREES/SHRUBS					
greasewood	Sarcobatus vermiculatus	SAVE4	8	190 - 380	5 - 10
rubber rabbitbrush	Ericameria nauseosa	ERNA10	9	0 - 190	0 - 5
short-fruit willow	Salix brachycarpa	SABR	10	0 - 190	0 - 5
Shrubby cinquefoil	Dasiphora floribunda	DAFL3	11	0 - 190	0 - 5

This list of plants and their relative proportions are based on near normal years. Fluctuations in species composition and relative production may change from year to year dependent upon precipitation or other climatic factors.

Plant Community Narratives

Following are the narratives for each of the described plant communities. These plant communities may not represent every possibility, but they probably are the most prevalent and repeatable plant communities. The plant composition tables shown above have been developed from the best available knowledge at the time of this revision. As more data is collected, some of these plant communities may be revised or removed, and new ones may be added. None of these plant communities should necessarily be thought of as "Desired Plant Communities". According to the USDA NRCS National Range and Pasture Handbook, Desired Plant Communities (DPC's) will be determined by the decision-makers and will meet minimum quality criteria established by the NRCS. The main purpose for including any description of a plant community here is to capture the current knowledge and experience at the time of this revision.

Alkali Sacaton/Basin Wildrye Plant Community (HCPC)

The interpretive plant community for this site is the Historic Climax Plant Community. This state evolved with grazing by large herbivores and is suited for grazing by domestic livestock. Potential vegetation is estimated at 75% grasses or grass-like plants, 10% forbs and 15% woody plants. Saline tolerant species dominate the site. The major grasses include alkali sacaton, basin wildrye, Nuttall's alkaligrass, alkali bluegrass, and inland saltgrass. Common woody plants are greasewood and rubber rabbitbrush.

A typical plant composition for this state consists of Alkali sacaton 40-50%, Basin wildrye 5-10%, Nuttall's alkaligrass 5-10%, alkali bluegrass 5-10%, up to 10% inland saltgrass, perennial forbs 1-10%, greasewood 5-10%, and up to 5% rubber rabbitbrush. Ground cover, by ocular estimate, varies from 70-80%.

The total annual production (air-dry weight) of this state is about 3800 pounds per acre, but it can range from about 3000 lbs./acre in unfavorable years to about 4500 lbs./acre in above average years.

The following is the growth curve of this plant community expected during a normal year:

Growth curve number: WY0203

Growth curve name: 15-19W, FREE WATER SITES

Growth curve description: WL, SB, SS FREE WATER SITES

JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC
0	0	0	0	10	40	30	15	5	0	0	0

(Monthly percentages of total annual growth)

The state is stable and well adapted to the Central Rocky Mountains climatic conditions. It is a critical state providing water and habitat for the surrounding area. It is resistant to drought due to a dependable water supply. This is a sustainable plant community (site/soil stability, watershed function, and biologic integrity).

Transitions or pathways leading to other plant communities are as follows:

- Continuous Season-long Grazing will convert this plant community to the *Alkali Muhly/Alkali Bluegrass State*.

Alkali Muhly/Alkali Bluegrass Plant Community

This plant community evolved under continuous grazing by domestic livestock. Saline tolerant grasses make up the majority of the understory. Dominant grasses include alkali muhly and alkali bluegrass.

The total annual production (air-dry weight) of this state is about 2800 pounds per acre, but it can range from about 2000 lbs./acre in unfavorable years to about 3500 lbs./acre in above average years.

The following is the growth curve of this plant community expected during a normal year:

Growth curve number: WY0203

Growth curve name: 15-19W, FREE WATER SITES

Growth curve description: WL, SB, SS FREE WATER SITES

JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC
0	0	0	0	10	40	30	15	5	0	0	0

(Monthly percentages of total annual growth)

The state is stable and protected from excessive erosion. The biotic integrity of this plant community is at risk due to decreased species diversity. The watershed is usually functioning.

Transitional pathways leading to other plant communities are as follows:

- Prescribed Grazing will result in a plant community very similar to the *Historic Climax Plant Community (Alkali Sacaton/Basin Wildrye State)*.

Ecological Site Interpretations

Animal Community – Wildlife Interpretations

Alkali Sacaton/Basin Wildrye Plant Community (HCPC): This plant community is very important for many of the wildlife species in the area. With the presence of water at or near the soil surface, over 80% of all wildlife will use this site to fulfill some part of their habitat needs. It provides forage for mule deer and antelope. It provides nesting habitat for shorebirds, songbirds, and waterfowl as well as ground nesting birds such as harriers. The lush herbaceous material produces insects for sage grouse brood rearing and foraging. Dense ground cover provides escape cover, forage, and breeding areas for small mammals which draw predators such as raptors, red fox and coyote. Other birds that would frequent this plant community include red-wing blackbirds, sandhill cranes, western meadowlarks, and neo-tropical migrants.

Alkali Muhly/Alkali Bluegrass Plant Community: This plant community may be beneficial for the same wildlife that would use the Historic Climax Plant Community. However, the plant community composition is less diverse, and thus, less apt to meet the seasonal needs of these animals.

Animal Preferences (Quarterly - 1,2,3,4) for commonly occurring plants in MLRA 43B, 15-19W

COMMON NAME/GROUP NAME	SCIENTIFIC NAME	SYMBOL	Cattle	Sheep	Horses	Mule Deer	Antelope	Elk	Moose
GRASSES/GRASSLIKES									
Alkali bluegrass	<i>Poa juncea</i> (syn. to <i>P. secunda</i>)	POJU	DDDD	PPPP	DDDD	PPPP	PPPP	DDDD	DDDD
Alkali muhly	<i>Muhlenbergia asperifolia</i>	MUAS	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD
Alkali sacaton	<i>Sporobolus airoides</i>	SPAI	PPPP	DDDD	PPPP	DDDD	DDDD	PPPP	DDDD
Alpine timothy	<i>Phleum alpinum</i>	PHAL2	PPPP	PPPP	PPPP	DDDD	UUUU	PPPP	DDDD
American manna grass	<i>Glyceria grandis</i>	GLGR	DDDD	UUUU	DDDD	UUUU	UUUU	DDDD	DDDD
Baltic rush	<i>Juncus balticus</i>	JUBA	DDDD	UUUU	DDDD	UUUU	UUUU	DDDD	UUUU
Basin wildrye	<i>Leymus cinereus</i>	LEC4	PPPP	PPPP	PPPP	DDDD	DDDD	PPPP	DDDD
Beaked sedge	<i>Carex rostrata</i>	CAR06	DDUD	UUUU	DDUD	UUUU	UUUU	DDUD	DDUD
Bearded wheatgrass	<i>Elymus trachycaulus</i> ssp. <i>subsecundus</i>	ELTRS	PPPP	DDDD	PPPP	DDDD	DDDD	PPPP	DDDD
Bentgrass	<i>Agrostis</i> spp.	AGROS2	PPPP	DDDD	PPPP	DDDD	DDDD	PPPP	DDDD
Big bluegrass	<i>Poa ampla</i> (syn. to <i>Poa secunda</i>)	POAM	PPPP	DDDD	PPPP	PPPP	PPPP	PPPP	PPPP
Blue wildrye	<i>Elymus glaucus</i>	ELGL	PPPP	DDDD	PPPP	DDDD	UUUU	PPPP	DDDD
Bluebunch wheatgrass	<i>Pseudoroegneria spicata</i>	FSSP6	PPPP	PPPP	PPPP	DDDD	DDDD	PPPP	DDDD
Bluejoint reedgrass	<i>Calamagrostis canadensis</i>	CACA4	PPPP	DDDD	PPPP	UUUU	UUUU	PPPP	DDDD
Bottlebrush squirreltail	<i>Elymus elymoides</i>	ELELE	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	NNNN
Bulrush	<i>Scirpus</i> spp.	SCIRP	DDDD	UUUU	UUUU	UUUU	UUUU	DDDD	DDDD
California oatgrass	<i>Danthonia californica</i>	DACA3	PPPP	DDDD	DDDD	DDDD	DDDD	PPPP	DDDD
Canby bluegrass	<i>Poa canbyi</i> (syn. to <i>Poa secunda</i>)	POCA	PPPP	DPDD	DPDD	DPDD	DPDD	PPPP	DPDD
Cattail	<i>Typha</i> spp.	TYPHA	DUUD	DUUD	DUUD	DUUD	DUUD	DUUD	DUUD
Columnia needlegrass	<i>Achnatherum nelsonii</i>	ACNE9	PPPP	DDDD	PPPP	DDDD	DDDD	PPPP	DDDD
Cusick bluegrass	<i>Poa cusickii</i>	POCU3	PPPP	PPPP	PPPP	PPPP	PPPP	PPPP	PPPP
Dunehead sedge	<i>Carex phaeocephala</i>	CAPH2	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Fowl bluegrass	<i>Poa palustris</i>	POPA2	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD
Green needlegrass	<i>Nassella viridula</i>	NAV14	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD
Idaho fescue	<i>Festuca idahoensis</i>	FEID	DDPD	DDPD	DDPD	DDDD	DDDD	DDPD	DDDD
Indian ricegrass	<i>Achnatherum hymenoides</i>	ACHY	PPPP	PPPP	PPPP	PPPP	PPPP	PPPP	PPPP
Inland saltgrass	<i>Distichlis spicata</i>	DISP	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Inland sedge	<i>Carex interior</i>	CAIN11	DDDD	DDDD	DDDD	UUUU	UUUU	DDDD	DDDD
Letterman needlegrass	<i>Achnatherum lettermanii</i>	ACLE9	UPUU	UPUU	UPUU	DDDD	DDDD	DDDD	DDDD
Little barley	<i>Hordeum pusillum</i>	HOPU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Mat muhly	<i>Muhlenbergia richardsonis</i>	MURI	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Montana wheatgrass	<i>Elymus albicans</i>	ELAL7	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD
Mountain brome	<i>Bromus marginatus</i>	BRMA4	PPPP	PPPP	DDDD	DDDD	NNNN	PPPP	DDDD
Mountain muhly	<i>Muhlenbergia montana</i>	MUMU	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD
Mutton bluegrass	<i>Poa fendleriana</i>	POFE	PPPP	PPPP	PPPP	PPPP	PPPP	PPPP	PPPP
Nebraska sedge	<i>Carex nebrascensis</i>	CANE2	PPPP	PPPP	PPPP	PPPP	DDDD	PPPP	DDDD
Needleleaf sedge	<i>Carex duriuscula</i>	CADU6	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Nodding brome	<i>Bromus porteri</i>	BRPO2	PPPP	PPPP	DDDD	DDDD	UUUU	PPPP	DDDD
Northern reedgrass	<i>Calamagrostis stricta</i> ssp. <i>inexpansa</i>	CAST13	PPPP	DDDD	PPPP	DDDD	UUUU	PPPP	DDDD
Nuttall's alkalgrass	<i>Puccinellia nuttalliana</i>	PUNU2	PPPP	PPPP	PPPP	PPPP	PPPP	PPPP	PPPP
One-spoke oatgrass	<i>Danthonia unispicata</i>	DAUN	DDDD	PPPP	DDDD	PPPP	DDDD	DDDD	DDDD
Oniongrass	<i>Melica bulbosa</i>	MEBU	PPPP	PPPP	PPPP	PPPP	PPPP	PPPP	PPPP
Prairie junegrass	<i>Koeleria macrantha</i>	KOMA	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD
Pumpelly's brome	<i>Bromus inermis</i> ssp. <i>pumpellianus</i>	BRINP	PPPP	PPPP	DDDD	DDDD	UUUU	PPPP	DDDD
Redtop	<i>Agrostis stolonifera</i>	AGST2	UPDU	UPDU	UPDU	UPDU	UPDU	UPDU	UPDU
Reed canarygrass	<i>Phalaris arundinacea</i>	PHAR3	UDDU	UDDU	UDDU	UDDU	UDDU	UDDU	UDDU
Richardson's needlegrass	<i>Achnatherum richardsonii</i>	ACHRI8	PPPP	PPPP	DDDD	DDDD	DDDD	PPPP	DDDD
Sandberg bluegrass	<i>Poa secunda</i>	POSE	UDDU	UDDU	UDDU	UDDU	UDDU	UDDU	UDDU
Shortawn foxtail	<i>Alopecurus aequalis</i>	ALAE	DDDU	DDDU	DDDU	DDDU	DDDU	DDDU	DDDU
Slender wheatgrass	<i>Elymus trachycaulus</i>	ELTR7	PPPP	DDDD	PPPP	DDDD	DDDD	PPPP	DDDD
Spikifescue	<i>Leucopoa kingii</i>	LEK12	PPPP	DDDD	PPPP	PPPP	DDDD	PPPP	DDDD
Spikerush	<i>Eleocharis</i> spp.	ELEOC	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Spike trisetum	<i>Trisetum spicatum</i>	TRSP2	PPPP	DDDD	PPPP	DDDD	DDDD	PPPP	DDDD
Sun sedge	<i>Carex inops</i> ssp. <i>heliophila</i>	CAINH2	PPPP	DDDD	PPPP	DDDD	DDDD	PPPP	DDDD
Tall manna grass	<i>Glyceria elata</i> (syn. <i>G. striata</i>)	GLEL	DDDD	UUUU	DDDD	UUUU	UUUU	DDDD	DDDD
Thickspike wheatgrass	<i>Elymus lanceolatus</i> ssp. <i>lanceolatus</i>	ELMA7	DPDD	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD
Threadleaf sedge	<i>Carex filifolia</i>	CAFI	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD
Timber oatgrass	<i>Danthonia intermedia</i>	DAIN	DDDD	DDDD	DDDD	UUUU	UUUU	DDDD	DDDD
Tufted hairgrass	<i>Deschampsia caespitosa</i>	DECA18	PPPP	PPPP	PPPP	DDDD	DDDD	PPPP	DDDD
Water sedge	<i>Carex aquatilis</i> ssp. <i>aquatilis</i>	CACA3	UDDU	UDDU	UDDU	UDDU	UDDU	UDDU	UDDU
Western needlegrass	<i>Achnatherum occidentale</i>	ACOOQ	PPPP	PPPP	PPPP	DDDD	DDDD	PPPP	DDDD
Western wheatgrass	<i>Pascopyrum smithii</i>	PASM	DPDD	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD
FORBS									
American licorice	<i>Glycyrrhiza lepidota</i>	GLLE3	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
American bistort	<i>Polygonum bistortoides</i>	POB16	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD
American vetch	<i>Vicia americana</i>	VIAM	PPPP	PPPP	PPPP	PPPP	PPPP	PPPP	DDDD
Arnica	<i>Arnica</i> spp.	ARNIC	UUUU	UUUU	UUUU	DDDD	UUUU	UUUU	UUUU
Arrowgrass	<i>Triglochin</i> spp.	TRIGL	TTTT	TTTT	TTTT	TTTT	TTTT	TTTT	TTTT
Asters	<i>Eucephalus & Symphyotrichum</i> spp.	EUCEP2/SYMPH4	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Avens (prairie smoke)	<i>Geum</i> spp.	GEUM	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Balsamorhiza	<i>Balsamorhiza</i> spp.	BALSA	DPDD	PPPP	PPPP	PPPP	PPPP	PPPP	PPPP
Bedstraw	<i>Galium</i> spp.	GALIU	UUUU	DDDD	UUUU	DDDD	DDDD	DDDD	UUUU
Biscuitroot	<i>Lomatium</i> spp.	LOMAT	DDDD	DDDD	UUUU	DDDD	DDDD	DDDD	DDDD
Bitterroot	<i>Lewisia rediviva</i> ssp. <i>rediviva</i>	LERER	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Bluebell	<i>Mertensia</i> spp.	MERTE	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD
Blue-eyed grass	<i>Sisyrinchium</i> spp.	SISYR	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Buckwheat	<i>Eriogonum</i> spp.	ERIOG	UUUU	DDDD	UUUU	UUUU	UUUU	UUUU	UUUU
Buttercup	<i>Ranunculus</i> spp.	RANUN	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD
Cinquefoil (herbaceous)	<i>Potentilla</i> spp.	POTEN	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Clover	<i>Trifolium</i> spp.	TRIFO	PPPP	PPPP	PPPP	PPPP	PPPP	PPPP	PPPP
Columbine	<i>Aquilegia</i> spp.	AQUIL	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD
Cow parsnip	<i>Hieracium maximum</i>	HERAC	PPPP	PPPP	PPPP	PPPP	PPPP	DDDD	NNNN
Daisy	<i>Townsendia</i> spp.	TOWNS	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Deathcamas	<i>Zigadenus venenosus</i>	ZIVE	TTTT	TTTT	TTTT	TTTT	TTTT	TTTT	TTTT
Elephanthead lousewort	<i>Pedicularis groenlandica</i>	PEGR2	UUUU	DDDD	UUUU	DDDD	UUUU	UUUU	DDDD
Elk thistle	<i>Cirsium foliosum</i>	CIFO	UDDU	UUUU	UDDU	UDDU	UUUU	UDDU	UUUU
Evening-primrose	<i>Oenothera</i> spp.	OENOT	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Fireweed	<i>Chamerion angustifolium</i>	CHAN9	PPPP	DDDD	UUUU	PPPP	DDDD	PPPP	PPPP
Flax	<i>Linum</i> spp.	LINUM	UPDU	UPDU	UPDU	UPDU	UPDU	UPDU	UPDU
Fleabane	<i>Erigeron</i> spp.	ERIGE2	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD
Gentian	<i>Gentiana</i> spp.	GENTI	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Geranium	<i>Geranium</i> spp.	GERAN	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Gilia	<i>Gilia</i> spp.	GILIA	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Goldenaster	<i>Heterotheca</i> spp.	HETER8	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Goldenpea	<i>Thermopsis</i> spp.	THERM	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Goldenrod	<i>Solidago</i> spp.	SOLID	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Golden smoke	<i>Corydalis aurea</i>	COAU2	TTUU	TTUU	TTUU	TTUU	TTUU	TTUU	TTUU
Goldenweed, stemless	<i>Stenotus acualis</i> ssp. <i>acualis</i>	STACA	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Green gentian	<i>Frasera speciosa</i>	FRSP	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD
Groundsel	<i>Packera</i> spp.	PACKE	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Harebell (bellflower)	<i>Campanula</i> spp.	CAMPA	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Hawksbeard	<i>Crepis</i> spp.	CREPI	UUUU	PPPP	UUUU	DDDD	DDDD	UUUU	DDDD
Hawkweed	<i>Hieracium</i> spp.	HIERA	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Horsemint	<i>Agastache</i> spp.	AGAST	DDDD	DDDD	UUUU	DDDD	DDDD	DDDD	DDDD
Horsetail (scouring rush)	<i>Equisetum</i> spp.	EQUIS	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Iris (Rocky Mountain)	<i>Iris missouriensis</i>	IRMI	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU

Animal Preferences (Quarterly - 1,2,3,4) for commonly occurring plants in MLRA 43B, 15-19W

COMMON NAME/GROUP NAME	SCIENTIFIC NAME	SYMBOL	Cattle	Sheep	Horses	Mule Deer	Antelope	Elk	Moose
Larkspur (poisonous in spring before flowering)	Delphinium spp.	DELPH	DTDD	DTDD	DTDD	DTDD	DTDD	DTDD	DTDD
Little sunflower	Helianthus pumilus	HEPU3	PPPP	PPPP	PPPP	PPPP	PPPP	PPPP	PPPP
Locoweed	Oxytropis spp.	LOXTR	TTUU	TTUU	TTUU	TTUU	TTUU	TTUU	TTUU
Lupine (may be poisonous after seedpots mature)	Lupinus spp.	LUPIN	DDTT	DDTT	DDTT	DDTT	DDTT	DDTT	DDTT
Meadow-rue	Thalictrum occidentale	THOC	DDDD	PPPP	DDDD	PPPP	PPPP	DDDD	PPPP
Milkvetch	Astragalus spp.	ASTRA	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD
Minerscandle	Cryptantha spp.	CRYPT	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Mint (wild)	Mentha arvensis	MEAR4	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Monkeyflower	Mimulus spp.	MIMUL	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Monkshood	Aconitum spp.	ACONI	TTTT	TTTT	TTTT	TTTT	TTTT	TTTT	TTTT
Mountain dandelion	Agoseris spp.	AGOSE	DDDD	PPPP	DDDD	PPPP	DDDD	DDDD	DDDD
Mule-ears	Wyethia amplexicaulis	WYAM	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Mustard	Draba spp.	DRABA	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Nailwort	Paronychia spp.	PARON	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Onion (wild)	Allium spp.	ALLI4	DPDD	PPPP	DPDD	DPDD	PPPP	DPDD	DPDD
Oregon grape	Mahonia repens	MARE11	UUUU	DDDD	UUUU	PPPP	DDDD	DDDD	DDDD
Owl's-clover	Orthocarpus spp.	ORTHO	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Paintbrush	Castilleja spp.	CAST	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD
Peavine	Lathyrus spp.	LATHY	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD
Penstemon	Penstemon spp.	PENST	PPPP	PPPP	PPPP	PPPP	PPPP	PPPP	PPPP
Phacelia	Phacelia spp.	PHACE	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD
Phlox	Phlox spp.	PHLOX	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Plantain	Plantago spp.	PLANT	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Primrose	Primula spp.	PRIMU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Pussytoes	Antennaria spp.	ANTEN	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Ragwort (groundsel)	Senecio spp.	SENEC	TTTT	TTTT	TTTT	TTTT	TTTT	TTTT	TTTT
Sandwort	Arenaria spp.	ARENA	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Shooting star	Dodecatheon spp.	DODEC	DDDD	DDDD	UUUU	DDDD	UUUU	UUUU	UUUU
Starwort	Stellaria spp.	STELL	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Sego lily	Calochortus nuttallii	CANU3	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Smartweed (knotweed)	Polygonum spp.	POLYG4	UUUU	UUUU	UUUU	DDDD	UUUU	UUUU	UUUU
Sneezeweed, orange (rubberweed)	Hymenoxys spp.	HYMEN7	TTTT	TTTT	TTTT	TTTT	TTTT	TTTT	TTTT
Springbeauty	Claytonia spp.	CLAYT	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Stinging nettle	Urtica dioica	URDI	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Stonecrop	Sedum spp.	SEDUM	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Stoneseed	Lithospermum spp.	LITHO3	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Sunflower	Helianthus spp.	HELIA3	PPPP	PPPP	PPPP	PPPP	PPPP	PPPP	PPPP
Sweetroot	Osmorhiza spp.	OSMOR	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD
Toadflax	Comandra umbellata	COUMP	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Valerian (tobacco root)	Valeriana spp.	VALER	DDDD	PPPP	DDDD	DDDD	DDDD	DDDD	DDDD
Violet	Viola spp.	VIOLA	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD
Water hemlock (spotted)	Cicuta maculata var. angustifolia	CIMAA	TTUU	TTUU	TTUU	TTUU	TTUU	TTUU	TTUU
Waterleaf	Hydrophyllum spp.	HYDR04	DDDD	DDDD	DDDD	PPPP	DDDD	DDDD	DDDD
Western coneflower	Rudbeckia occidentalis	RUOC2	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Wild strawberry (false strawberry)	Fragaria vesca	FRVE	DDDD	PPPP	DDDD	PPPP	PPPP	DDDD	DDDD
Yarrow (common & western)	Achillea millefolium	ACMI2	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Yellowbell	Fritillaria pudica	FRPU2	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Yellow sneezeweed	Helenium autumnale	HEAU	TTTT	TTTT	TTTT	TTTT	TTTT	TTTT	TTTT
TREES, SHRUBS & HALF-SHRUBS									
Alpine laurel (bog kalmia)	Kalmia microphylla	KAMI	TTTT	TTTT	TTTT	TTTT	TTTT	TTTT	TTTT
Antelope bitterbrush	Purshia tridentata	PUTR2	PPPP	PPPP	DDDD	PPPP	PPPP	PPPP	PPPP
Aspen	Populus tremuloides	POTR5	DDDD	DDDD	DDDD	PPPP	DDDD	PPPP	PPPP
Basin big sagebrush	Artemisia tridentata ssp. tridentata	ARTR1	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Big sagebrush	Artemisia tridentata	ARTR2	UUUU	DDDD	UUUU	PPPP	PPPP	DDDD	DDDD
Black sagebrush	Artemisia nova	ARN04	DDDD	PPPP	UUUU	DDDD	DDDD	DDDD	DDDD
Chokecherry (toxic in large amounts)	Prunus virginiana	PRVI	DTDD	DTDD	DDDD	PPPP	UUUU	DDDD	PPPP
Current	Ribes spp.	RIBES	DDDD	DDDD	DDDD	PPPP	UUUU	DDDD	DDDD
Dogwood	Cornus spp.	CORNU	DDDD	DPDD	DDDD	DPDD	DDDD	DPDD	DPDD
Early (alkali) sage	Artemisia arbuscula ssp. longiloba	ARALR	UUUU	UUUU	UUUU	PPPP	PPPP	UUUU	UUUU
Elderberry	Sambucus spp.	SAMBU	DDDD	DDDD	UUUU	PPPP	UUUU	DDDD	DDDD
Fringed sagewort	Artemisia frigida	ARFR4	UUUU	UUUU	UUUU	UUUU	DDDD	UUUU	UUUU
Goldenweed, shrubby	Ericameria suffruticosa	ERSU13	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Greasewood (toxic in large amounts)	Sarcobatus vermiculatus	SAVE4	UUUU	DDDD	UUUU	DDDD	DDDD	UUUU	UUUU
Green (low) rabbitbrush	Chrysothamnus viscidiflorus	CHV18	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Juniper, common	Juniperus communis var. depressa	JUCOD	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Juniper, Rocky Mountain	Juniperus scopulorum	JUSC2	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Limber pine	Pinus flexilis	PIFL2	NNNN	NNNN	NNNN	NNNN	NNNN	NNNN	NNNN
Low sagebrush	Artemisia arbuscula ssp. arbuscula	ARAR8	UUUU	DDDD	UUUU	DDDD	DDDD	UUUU	UUUU
Mountain big sagebrush	Artemisia tridentata ssp. vaseyana	ARTRV	UUUU	DDDD	UUUU	DDDD	DDDD	UUUU	UUUU
Mountain mahogany	Cercocarpus spp.	CERCO	PPPP	PPPP	DDDD	PPPP	UUUU	PPPP	PPPP
Raspberry	Rubus idaeus	RUID	UUUU	UUUU	UUUU	DDDD	UUUU	UUUU	DDDD
Rubber rabbitbrush	Ericameria nauseosa	ERNA10	UUUU	PPPP	UUUU	DDDD	PPPP	DDDD	DDDD
Serviceberry	Amelanchier alnifolia	AMAL2	DDDD	PPPP	UUUU	PPPP	DDDD	DDDD	DDDD
Shrubby cinquefoil	Dasiphora floribunda	DAFL3	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Silverberry	Elaeagnus commutata	ELCO	UUUU	DDDD	UUUU	UUUU	DDDD	UUUU	PPPP
Silver sagebrush	Artemisia cana	ARCA13	UUUU	DDDD	UUUU	PPPP	PPPP	DDDD	DDDD
Snowberry (western)	Symphoricarpos occidentalis	SYOC	UUUU	UUUU	UUUU	DDDD	UUUU	UUUU	UUUU
Snowbrush ceanothus	Ceanothus velutinus	CEVE	UUUU	DDDD	UUUU	DDDD	UUUU	DDDD	UUUU
Spiked big sagebrush	Artemisia tridentata ssp. spiciformis	ARTRS2	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Thimbleberry	Rubus parviflorus	RUPA	UUUU	DDDD	UUUU	UUUU	UUUU	UUUU	DPDD
Three-tip sagebrush	Artemisia tripartita	ARTR4	UUUU	DDDD	UUUU	UUUU	DDDD	UUUU	DDDD
True mountainmahogany	Cercocarpus montanus	CEMO2	PPPP	PPPP	DDDD	PPPP	UUUU	PPPP	PPPP
Water birch	Betula occidentalis	BEOC2	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD
Wild rose	Rosa woodsii var. woodsii	ROWOW	DDDD	DDDD	UUUU	DDDD	DDDD	DDDD	DDDD
Willow, Bebb's	Salix bebbiana	SABE2	DDDD	PPPP	DDDD	PPPP	DDDD	DDDD	PPPP
Willow, Blueberry	Salix myrtillofolia	SAMY	DDDD	PPPP	DDDD	PPPP	DDDD	DDDD	PPPP
Willow, Booth's	Salix boothii	SABO2	DDDD	PPPP	DDDD	PPPP	DDDD	DDDD	PPPP
Willow, coyote (sandbar)	Salix exigua	SAEX	PPPP	PPPP	DDDD	PPPP	UUUU	PPPP	PPPP
Willow, Drummonds	Salix drummondiana	SADR	DDDD	PPPP	DDDD	PPPP	DDDD	DDDD	PPPP
Willow, grayleaf	Salix glauca	SAGL	DDDD	PPPP	DDDD	PPPP	DDDD	DDDD	PPPP
Willow, Geyers	Salix geyeriana	SAGE2	DDDD	PPPP	DDDD	PPPP	DDDD	DDDD	PPPP
Willow, Lemmons	Salix lemmonii	SALE	DDDD	PPPP	DDDD	PPPP	DDDD	DDDD	PPPP
Willow, peachleaf	Salix amygdaloides	SAAM2	PPPP	PPPP	DDDD	PPPP	UUUU	PPPP	PPPP
Willow, planeleaf (diamondleaf)	Salix planifolia	SAPL2	DDDD	PPPP	DDDD	PPPP	DDDD	DDDD	PPPP
Willow, pussy	Salix discolor	SADI	DDDD	DDDD	DDDD	DDDD	UUUU	DDDD	DDDD
Willow, Scoulers	Salix scouleriana	SASC	PPPP	PPPP	DDDD	PPPP	DDDD	PPPP	PPPP
Willow, short-fruit (barrenground)	Salix brachycarpa	SABR	DDDD	PPPP	DDDD	PPPP	DDDD	DDDD	PPPP
Willow, tweedy	Salix tweedyi	SATW	DDDD	PPPP	DDDD	PPPP	DDDD	DDDD	PPPP
Willow, whiplash	Salix lucida ssp. Caudata	SALUC	DDDD	PPPP	DDDD	PPPP	DDDD	DDDD	PPPP
Willow, interior	Salix interior	SAIN3	DDDD	PPPP	DDDD	PPPP	DDDD	DDDD	PPPP
Willow, wolf	Salix woffii	SAWO	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	DDDD
Willow, yellow	Salix lutea	SALU2	PPPP	PPPP	DDDD	PPPP	UUUU	PPPP	PPPP
Wyoming big sagebrush	Artemisia tridentata ssp. wyomingensis	ARTRW8	UUUU	DDDD	UUUU	PPPP	PPPP	UUUU	UUUU

N = not used; U = undesirable; D = desirable; P = preferred; T = toxic

Animal Community – Grazing Interpretations

The following table lists suggested stocking rates for cattle under continuous season-long grazing under normal growing conditions. These are conservative estimates that should be used only as guidelines in the initial stages of the conservation planning process. Often, the current plant composition does not entirely match any particular plant community (as described in this ecological site description). Because of this, a field visit is recommended, in all cases, to document plant composition and production. More precise carrying capacity estimates should eventually be calculated using this information along with animal preference data, particularly when grazers other than cattle are involved. Under more intensive grazing management, improved harvest efficiencies can result in an increased carrying capacity. If distribution problems occur, stocking rates must be reduced to maintain plant health and vigor.

Plant Community	Production (lb./ac)	Carrying Capacity* (AUM/ac)
Alkali Sacaton/Basin Wildrye (HCPC)	3000-4500	1.2
Alkali Muhly/Alkali Bluegrass	2000-3500	.9

* - Continuous, season-long grazing by cattle under average growing conditions.

Grazing by domestic livestock is one of the major income-producing industries in the area. Rangeland in this area may provide yearlong forage for cattle, sheep, or horses. During the dormant period, the forage for livestock use needs to be supplemented with protein because the quality does not meet minimum livestock requirements.

Hydrology Functions

Salinity/alkalinity is the principal factor limiting forage production on this site. This site is dominated by soils in hydrologic group C and D. Infiltration ranges from moderately slow to moderately well. Runoff potential for this site varies from moderate to high depending on soil hydrologic group and ground cover. In many cases, areas with greater than 75% ground cover have the greatest potential for high infiltration and lower runoff. Areas where ground cover is less than 50% have the greatest potential to have reduced infiltration and higher runoff (refer to Part 630, NRCS National Engineering Handbook for detailed hydrology information).

Rills and gullies should not typically be present. Water flow patterns should be barely distinguishable if at all present. Pedestals are only slightly present in association with bunchgrasses. Litter typically falls in place, and signs of movement are not common. Chemical and physical crusts are sometimes present.

Recreational Uses

This site provides a variety of hunting opportunities as well as providing popular camping areas for recreationists. This site has a wide variety of forbs which bloom throughout spring and summer, providing esthetic values that appeal to visitors.

Wood Products

No appreciable wood products are present on the site.

Other Products

Supporting Information

Associated Sites

Overflow	R043BY230WY
Subirrigated	R043BY274WY
Wetland	R043BY278WY

Similar Sites

R034AY242WY – Saline Subirrigated (SS) 10-14W has lower production.

R034AY238WY – Saline Lowland (SL) 10-14W has lower production and is dominated by greasewood.

Inventory Data References (narrative)

Information presented here has been derived from NRCS clipping data and other inventory data. Field observations from range trained personnel were also used. Those involved in developing this site include: Bill Christensen, Range Management Specialist, NRCS; Karen Clause, Range Management Specialist, NRCS; and Everet Bainter, Range Management Specialist, NRCS. Other sources used as references include: USDA NRCS Water and Climate Center, USDA NRCS National Range and Pasture Handbook, and USDA NRCS Soil Surveys from various counties.

Inventory Data References

<u>Data Source</u>	<u>Number of Records</u>	<u>Sample Period</u>	<u>State</u>	<u>County</u>
SCS-RANGE-417	58	1966-1986	WY	Lincoln & others

State Correlation

Type Locality

Field Offices

Lyman, Cokeville, Afton, Jackson, Pinedale

Relationship to Other Established Classifications

Other References

Site Description Approval

State Range Management Specialist

Date